

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : A61K 47/14, A61L 15/44	A1	(11) International Publication Number: <b>WO 96/37231</b> (43) International Publication Date: 28 November 1996 (28.11.96)
---	----	---

(21) International Application Number: PCT/US96/06868

(22) International Filing Date: 13 May 1996 (13.05.96)

(30) Priority Data:  
08/452,468 26 May 1995 (26.05.95) US

(71) Applicant: ALZA CORPORATION [US/US]; 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US).


(72) Inventors: LEE, Eun, Soo; 108 West Danbury, Redwood City, CA 94061 (US). YUM, Su, II; 1021 Runnymede Court, Los Altos, CA 94021 (US).

(74) Agents: RAFA, Michael, J. et al.; Alza Corporation, 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US).

(81) Designated States: AU, CA, CN, JP, KR, MX, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

**Published**  
*With international search report.*

(54) Title: SKIN PERMEATION ENHANCER COMPOSITIONS USING ACYL LACTYLATES



## (57) Abstract

The present invention is directed to the transdermal administration of a drug together with a suitable amount of an acyl lactylate permeation enhancer. The invention includes a transdermal drug delivery device comprising a matrix adapted to be placed in drug-and permeation enhancer-transmitting relation with a skin site. The matrix contains sufficient amounts of an acyl lactylate permeation enhancer and drug, in combination, to continuously administer drug to the systemic circulation of a patient at a therapeutically effective rate. The invention is also directed to compositions and methods for transdermal administration of a drug together with an acyl lactylate permeation enhancer, alone or in combination with other enhancers.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

# **SKIN PERMEATION ENHANCER COMPOSITIONS**

## **USING ACYL LACTYLATES**

### **FIELD OF THE INVENTION**

This invention relates to the transdermal delivery of drugs or other biologically active agents and more particularly to methods and compositions for enhancing the percutaneous absorption of drugs or other agents when incorporated in transdermal drug delivery systems or devices. Particularly, this invention relates to the use of acyl lactylates as permeation enhancers for transdermal systems or compositions.

### **DESCRIPTION OF TERMS**

As used herein, the term "transdermal" delivery or administration refers to the delivery or administration of agents by permeation through a body surface or membrane, preferably intact skin, by topical administration.

As used herein, the term "therapeutically effective" amount or rate refers to the amount or rate of drug or active agent needed to effect the desired therapeutic result.

As used herein, the phrase "sustained time period" intends at least about 12 hours and will typically intend a period in the range of about one to about seven days.

As used herein, the phrase "predetermined area of skin" intends a defined area of intact unbroken skin or mucosal tissue. That area will usually be in the range of about 5 cm<sup>2</sup> to about 100 cm<sup>2</sup>.

1           As used herein, the expressions "drug" and "agent" are used  
2 interchangeably and are intended to have their broadest interpretation as to  
3 any therapeutically active substance which is delivered to a living organism to  
4 produce a desired, usually beneficial, effect.

## 5 6                                   BACKGROUND OF THE INVENTION

7  
8           Acyl lactylates are represented by the general structure:  
9  $R-CO-(OCH_2CH_2CO)_n-OH$  where R is a straight or branched alkyl or aryl  
10 group consisting of 3 to 20 carbons and  $n = 1$  to 10. Acyl lactylates have  
11 been used in the food industry as dough conditioners and softeners and as  
12 oil-in-water emulsifiers in nondairy compositions such as coffee whiteners and  
13 vegetable oil based whipped toppings. They have also been used as  
14 emulsifiers in analgesic stick compositions and in conjunction with other  
15 co-emulsifiers to produce an emulsion base for cosmetic or pharmaceutical  
16 compositions. Typically, acyl lactylates are commercially available as a salt  
17 form for use in cosmetic formulations. The salt form is not effective as a  
18 permeation enhancer.

19           Other uses of acyl lactylates include their use as antimicrobial or  
20 antibacterial agents, and as a protectant against hair loss or in topical  
21 compositions for inducing, maintaining, or increasing hair growth. In general,  
22 permeation enhancers that are not normally toxic at the concentrations  
23 employed in cosmetic or medical compositions may exhibit toxic effects at the  
24 higher concentrations required to produce adequate permeation  
25 enhancement. Accordingly, acyl lactylates have not been used as a  
26 permeation enhancer alone or in combination with other permeation  
27 enhancing agents in order to increase the permeability of a body surface or  
28 membrane to a drug or active agent to transdermally deliver the drug or agent  
29 to the systemic circulation of a patient.

1           US Patent No. 4,184,978, incorporated herein in its entirety by  
2 reference, describes stable oil-in-water emulsion systems for use in  
3 cosmetics, toiletries, and pharmaceuticals. Acyl lactylates are disclosed as  
4 suitable emulsifiers.

5           US Patent No. 4,301,820, incorporated herein in its entirety by  
6 reference, describes a composition comprising at least one fatty acid lactylate  
7 and/or glycolate as a humectant compound in permanent waving  
8 compositions.

9           U.S. Patent No. 4,702,916, incorporated herein in its entirety by  
10 reference, describes the use of acyl lactylates as emulsifiers in analgesic gel  
11 stick compositions.

12           U.S. Patent No. 5,124,354, incorporated herein in its entirety by  
13 reference, describes a special protein tyrosine kinase inhibitor and a  
14 cosmetically acceptable vehicle. Surface active agents, including acyl  
15 lactylates, are disclosed as penetration enhancers that improve delivery of the  
16 composition to its site of action in the immediate environment of the hair  
17 follicle close to the dermal papilla.

18           European Patent Application 0 573 253 incorporated herein in its  
19 entirety by reference describes an anti-bacterial cosmetic composition for  
20 topical application to the skin and/or hair comprising C<sub>6</sub>-C<sub>12</sub> acyl lactylate or a  
21 derivative thereof as an anti-bacterial substance. The composition is  
22 especially beneficial in the treatment of unwanted hair loss.

23           European Patent Application 0 572 271 incorporated herein in its  
24 entirety by reference describes the use of acyl lactylates as preservatives in  
25 topical cosmetic compositions for prevention of the growth of undesired  
26 microorganisms in the skin or hair.

27           The transdermal route of parenteral delivery of drugs provides many  
28 advantages, and transdermal systems for delivering a wide variety of drugs or  
29 other beneficial agents are described in U.S. Pat. Nos. 3,598,122; 3,598,123;  
30 3,731,683; 3,797,494; 4,286,592; 4,314,557; 4,379,454; 4,435,180;

1 4,559,222; 4,568,343; 4,573,999; 4,588,580; 4,645,502; 4,704,282;  
2 4,816,258; 4,849,226; 4,908,027; 4,943,435; and 5,004,610, for example,  
3 all of which are incorporated herein by reference. In many cases, drugs  
4 which would appear to be ideal candidates for transdermal delivery are found  
5 to have such low permeability through intact skin that they cannot be  
6 delivered in therapeutically effective amounts from reasonably sized devices.

7 In an effort to increase skin permeability so that drugs can be delivered  
8 in therapeutically effective amounts, it has been proposed to pretreat the skin  
9 with various chemicals or to concurrently deliver the drug in the presence of a  
10 permeation enhancer. Various materials have been suggested for this,  
11 as described in U.S. Patent Nos. 3,472,931; 3,527,864; 3,896,238;  
12 3,903,256; 3,952,099; 4,046,886; 4,130,643; 4,130,667; 4,299,826;  
13 4,335,115; 4,343,798; 4,379,454; 4,405,616; 4,746,515; 4,788,062;  
14 4,820,720; 4,863,738; 4,863,970; and 5,378,730; British Patent  
15 No. 1,011,949; and Idson, "Percutaneous Absorption,"  
16 J. Pharm. Sci. (1975) 64:901-924.

17 To be considered useful, a permeation enhancer should have the  
18 ability to enhance the permeability of the skin for at least one and preferably a  
19 significant number of drugs. More importantly, it should be able to enhance  
20 the skin permeability such that the drug delivery rate from a reasonably sized  
21 system (preferably 5-50cm<sup>2</sup>) is at therapeutically effective levels. Additionally,  
22 the enhancer when applied to the skin surface, should be non-toxic, non-  
23 irritating on prolonged exposure and under occlusion, and non-sensitizing on  
24 repeated exposure. Preferably, it should be odorless and capable of  
25 delivering drugs without producing burning or tingling sensations.

## SUMMARY OF THE INVENTION

According to the present invention, it has been discovered that acyl lactylates are effective in enhancing the permeation of drugs through body surfaces and membranes generally, and through skin in particular. Importantly, the acyl lactylates of the present invention are able to enhance the permeability of drugs such that they can be delivered at therapeutically effective rates with reasonably sized transdermal delivery devices. Preferred acyl lactylate permeation enhancers of the present invention are caproyl lactic acid and lauroyl lactic acid.

Accordingly, the present invention provides a composition of matter for administration to a body surface or membrane to deliver at least one drug to the systemic circulation of a patient, at a therapeutically effective rate, by permeation through the body surface or membrane, comprising at least one drug and a permeation-enhancing amount of an acyl lactylate. The invention further provides a method for the transdermal coadministration of a drug at a therapeutically effective rate together with a skin permeation-enhancing amount of an acyl lactylate.

The system of the invention is preferably a transdermal drug delivery device comprising a matrix adapted to be placed in drug- and permeation enhancer-transmitting relation with the skin or mucosa. The system must be of a reasonable size useful for the application of the drug and the enhancer to a human body.

The invention is further directed to a composition of matter which optionally includes, in addition to the drug and acyl lactylate, one or more additional permeation enhancing compounds and to a method for the transdermal coadministration of such a composition.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a cross-sectional view of one embodiment of a transdermal therapeutic drug delivery device which may be used in accordance with the present invention.

FIG. 2 is a cross-sectional view of another embodiment of a transdermal therapeutic drug delivery device which may be used in accordance with the present invention.

FIG. 3 is a cross-sectional view of yet another embodiment of a transdermal therapeutic drug delivery device which may be used in accordance with this invention.

FIG. 4 is a graph of the flux of progesterone through human epidermis at 35° C, in vitro, from a mineral oil system with various permeation enhancers.

FIG. 5 is a graph of the flux of testosterone through human epidermis at 35° C, in vitro, from a mineral oil system with various permeation enhancers.

FIG. 6 is a graph of the flux of testosterone through human epidermis at 35° C, in vitro, from an EVA matrix system with various permeation enhancers.

## DETAILED DESCRIPTION OF THE INVENTION

According to this invention, it has been discovered that acyl lactylates can be used to effectively enhance the permeability of drugs through body surfaces or membranes and particularly through the skin. Specifically, it has been found that acyl lactylates, when converted to a free acid form, enhance the permeability of the body surface or membrane to drugs or other biologically active agents such that therapeutically effective amounts of a drug or agent can be systemically delivered from reasonably sized devices at

1 therapeutically effective rates. Additionally, it has been found that water  
2 dispersion of the acid form does not show corrosively low pH (pH 4.0-5.0).

3 It is believed that this invention has utility in connection with the  
4 delivery of drugs within the broad class normally delivered through body  
5 surfaces and membranes, including skin. In general, this includes therapeutic  
6 agents in all of the major areas, including, but not limited to, ACE inhibitors,  
7 adenylosuccinyl hormones, adrenergic neuron blocking agents,  
8 adrenocortical steroids, inhibitors of the biosynthesis of adrenocortical  
9 steroids, alpha-adrenergic agonists, alpha-adrenergic antagonists, selective  
10 alpha-two-adrenergic agonists, analgesics, antipyretics and anti-inflammatory  
11 agents, androgens, local and general anesthetics, antiaddictive agents,  
12 antiandrogens, antiarrhythmic agents, antiasthmatic agents, anticholinergic  
13 agents, anticholinesterase agents, anticoagulants, antidiabetic agents,  
14 antidiarrheal agents, antidiuretic, antiemetic and prokinetic agents,  
15 antiepileptic agents, antiestrogens, antifungal agents, antihypertensive  
16 agents, antimicrobial agents, antimigraine agents, antimuscarinic agents,  
17 antineoplastic agents, antiparasitic agents, antiparkinson's agents, antiplatelet  
18 agents, antiprogestins, antithyroid agents, antitussives, antiviral agents,  
19 atypical antidepressants, azaspirodecanediones, barbituates,  
20 benzodiazepines, benzothiadiazides, beta-adrenergic agonists, beta-  
21 adrenergic antagonists, selective beta-one-adrenergic antagonists, selective  
22 beta-two-adrenergic agonists, bile salts, agents affecting volume and  
23 composition of body fluids, butyrophenones, agents affecting calcification,  
24 calcium channel blockers, cardiovascular drugs, catecholamines and  
25 sympathomimetic drugs, cholinergic agonists, cholinesterase reactivators,  
26 dermatological agents, diphenylbutylpiperidines, diuretics, ergot alkaloids,  
27 estrogens, ganglionic blocking agents, ganglionic stimulating agents,  
28 hydantoins, agents for control of gastric acidity and treatment of peptic ulcers,  
29 hematopoietic agents, histamines, histamine antagonists,

1 5-hydroxytryptamine antagonists, drugs for the treatment of  
 2 hyperlipoproteinemia, hypnotics and sedatives, immunosuppressive agents,  
 3 laxatives, methylxanthines, monoamine oxidase inhibitors, neuromuscular  
 4 blocking agents, organic nitrates, opioid analgesics and antagonists,  
 5 pancreatic enzymes, phenothiazines, progestins, prostaglandins, agents for  
 6 the treatment of psychiatric disorders, retinoids, sodium channel blockers,  
 7 agents for spasticity and acute muscle spasms, succinimides, thioxanthines,  
 8 thrombolytic agents, thyroid agents, tricyclic antidepressants, inhibitors of  
 9 tubular transport of organic compounds, drugs affecting uterine motility,  
 10 vasodilators, vitamins and the like.

11 Representative drugs include, by way of example and not for purposes  
 12 of limitation, bepridil, diltiazem, felodipine, isradipine, nicardipine, nifedipine,  
 13 nimodipine, nitredipine, verapamil, dobutamine, isoproterenol, carterolol,  
 14 labetalol, levobunolol, nadolol, penbutolol, pindolol, propranolol, sotalol,  
 15 timolol, acebutolol, atenolol, betaxolol, esmolol, metoprolol, albuterol,  
 16 bitolterol, isoetharine, metaproterenol, pirbuterol, ritodrine, terbutaline,  
 17 alclometasone, aldosterone, amcinonide, beclomethasone, dipropionate,  
 18 betamethasone, clobetasol, clocortolone, cortisol, cortisone, corticosterone,  
 19 desonide, desoximetasone, 11-desoxycorticosterone, 11-desoxycortisol,  
 20 dexamethasone, diflorasone, fludrocortisone, flunisolide, fluocinolone,  
 21 fluocinonide, fluorometholone, flurandrenolide, halcinonide, hydrocortisone,  
 22 medrysone, 6 $\alpha$ -methylprednisolone, mometasone, paramethasone,  
 23 prednisolone, prednisone, tetrahydrocortisol, triamcinolone, benoxinate,  
 24 benzocaine, bupivacaine, chloroprocaine, cocaine, dibucaine, dyclonine,  
 25 etidocaine, lidocaine, mepivacaine, pramoxine, prilocaine, procaine,  
 26 proparacaine, tetracaine, alfentanil, chloroform, clonidine, cyclopropane,  
 27 desflurane, diethyl ether, droperidol, enflurane, etomidate, fentanyl,  
 28 halothane, isoflurane, ketamine hydrochloride, meperidine, methohexital,  
 29 methoxyflurane, morphine, propofol, sevoflurane, sufentanil, thiamylal,  
 30 thiopental, acetaminophen, allopurinol, apazone, aspirin, auranofin,

1 aurothioglucose, colchicine, diclofenac, diflunisal, etodolac, fenoprofen,  
 2 flurbiprofen, gold sodium thiomalate, ibuprofen, indomethacin, ketoprofen,  
 3 meclofenamate, mefenamic acid, meselamine, methyl salicylate,  
 4 nabumetone, naproxen, oxyphenbutazone, phenacetin, phenylbutazone,  
 5 piroxicam, salicylamide, salicylate, salicylic acid, salsalate, sulfasalazine,  
 6 sulindac, tolmetin, acetophenazine, chlorpromazine, fluphenazine,  
 7 mesoridazine, perphenazine, thioridazine, trifluorperazine, triflupromazine,  
 8 disopyramide, encainide, flecainide, indecainide, mexiletine, moricizine,  
 9 phenytoin, procainamide, propafenone, quinidine, tocainide, cisapride,  
 10 domperidone, dronabinol, haloperidol, metoclopramide, nabilone,  
 11 prochlorperazine, promethazine, thiethylperazine, trimethobenzamide,  
 12 buprenorphine, butorphanol, codeine, dezocine, diphenoxylate, drocode,  
 13 hydrocodone, hydromorphone, levallorphan, levorphanol, loperamide,  
 14 meptazinol, methadone, nalbuphine, nalmefene, nalorphine, naloxone,  
 15 naltrexone, oxybutynin, oxycodone, oxymorphone, pentazocine,  
 16 propoxyphene, isosorbide dinitrate, nitroglycerin, theophylline, phenylephrine,  
 17 ephedrine, pilocarpine, furosemide, tetracycline, chlorpheniramine, ketorolac,  
 18 bromocriptine, guanabenz, prazosin, doxazosin, and flufenamic acid.

19 Other representative drugs include benzodiazepines, such as  
 20 alprazolam, brotizolam, chlordiazepoxide, clobazam, clonazepam,  
 21 clorazepate, demoxepam, diazepam, flumazenil, flurazepam, halazepam,  
 22 lorazepam, midazolam, nitrazepam, nordazepam, oxazepam, prazepam,  
 23 quazepam, temazepam, triazolam, and the like; an antimuscarinic agent such  
 24 as anisotropine, atropine, clidinium, cyclopentolate, dicyclomine, flavoxate,  
 25 glycopyrrolate, hexocyclium, homatropine, ipratropium, isopropamide,  
 26 mepenzolate, methantheline, oxyphencyclimine, pirenzepine, propantheline,  
 27 scopolamine, telenzepine, tridihexethyl, tropicamide, and the like; an estrogen  
 28 such as chlorotrianisene, siethylstilbestrol, methyl estradiol, estrone, estrone  
 29 sodium sulfate, estropipate, mestranol, quinestrol, sodium equilin sulfate,  
 30 17 $\beta$ -estradiol (or estradiol), semi-synthetic estrogen derivatives such as the

1 esters of natural estrogen, such as estradiol-17 $\beta$ -enanthate, estradiol-17 $\beta$ -  
2 valerate, estradiol-3-benzoate, estradiol-17 $\beta$ -undecenoate, estradiol 16,  
3 17-hemisuccinate or estradiol-17 $\beta$ -cypionate, and the 17-alkylated estrogens,  
4 such as ethinyl estradiol, ethinyl estradiol-3- isopropylsulphonate, and the  
5 like; an androgen such as danazol, fluoxymesterone, methandrostenolone,  
6 methyltestosterone, nandrolone decanoate, nandrolone phenpropionate,  
7 oxandrolone, oxymetholone, stanozolol, testolactone, testosterone,  
8 testosterone cypionate, testosterone enanthate, testosterone propionate,  
9 and the like; or a progestin such as ethynodiol diacetate, gestodene,  
10 hydroxyprogesterone caproate, levonorgestrel, medroxyprogesterone  
11 acetate, megestrol acetate, norethindrone, norethindrone acetate,  
12 norethynodrel, norgestrel, progesterone, and the like.

13 Administration of the drug according to the invention comprises  
14 administering the drug at a therapeutically effective rate to an area of a body  
15 surface or membrane and simultaneously administering an acyl lactylate to  
16 the area of the body surface or membrane at a rate sufficient to substantially  
17 increase the permeability of the area to the drug formulation.

18 According to the invention, an acyl lactylate permeation enhancer and  
19 the drug to be delivered are placed in drug- and acyl lactylate-transmitting  
20 relationship to the appropriate body surface, preferably in a carrier therefor,  
21 and maintained in place for the desired period of time. The drug and acyl  
22 lactylate are typically dispersed within a physicochemically and biologically  
23 compatible matrix or carrier which may be applied directly to the body surface  
24 or skin as an ointment, gel, cream, suppository or sublingual or buccal tablet,  
25 for example, but are more preferably administered from a transdermal  
26 therapeutic delivery device as more fully described below. When used in the  
27 form of a liquid, ointment, cream, or gel applied directly to the skin, it is  
28 preferable, although not required, to occlude the site of administration.  
29 Such compositions can also contain other permeation enhancers, stabilizers,  
30 dyes, diluents, pigments, vehicles, inert fillers, excipients, gelling agents,

1 vasoconstrictors, and other components of typical compositions as are known  
2 to the art.

3 The acyl lactylates of this invention have a permeation-enhancing  
4 effect on the transport of drugs through body surface tissues generally,  
5 in addition to the skin. However, because skin is one of the most effective  
6 body barriers to the permeation of foreign substances, the effect of acyl  
7 lactylates on skin permeation makes it extremely useful in transdermal  
8 delivery. The following description of embodiments of the invention is  
9 therefore directed primarily to improving systemic delivery of these drugs by  
10 permeation through the skin.

11 It may be desirable in certain instances or with certain drugs to include  
12 one or more additional permeation enhancers in combination with the acyl  
13 lactylate. Thus, in certain embodiments of the present invention, a second  
14 permeation enhancer is included together with the drug and acyl lactylate  
15 permeation enhancer. This second enhancer may be selected from those  
16 compounds that have a permeation-enhancing effect with the drug and are  
17 compatible with the drug and with the acyl lactylate. For example, the second  
18 permeation enhancer may be a monoglyceride or mixture of monoglycerides  
19 of fatty acids such as glycerol monolaurate (GML) or glycerol monooleate  
20 (GMO), lauramide diethanolamine (LDEA), esters of fatty acids having from  
21 about 10 to 20 carbon atoms, and/or a lower C<sub>1-4</sub> alcohol such as ethanol or  
22 isopropanol.

23 Typically, monoglycerides have been available as a mixture of  
24 monoglycerides of fatty acids with one monoglyceride being the principal  
25 component, from which component the mixture derives its name.  
26 For example, one commercial monoglyceride is Emerest 2421 glycerol  
27 monooleate (Emery Division, Quanturn Chemical Corp.), which is a mixture of  
28 glycerol oleates with a glycerol monooleate content of 58% and a total  
29 monoesters content of 58%. Other examples of commercial monoglycerides  
30 are Myverol 1899K glycerol monooleate (Eastman Chemical Products) which

1 has a glycerol monooleate content of 61% and a total monoesters content of  
2 93%, and Myverol 1892K glycerol monolinoleate which has a glycerol  
3 monolinoleate content of 68% and a minimum total monoesters content of  
4 90%. The monoesters are chosen from those with from 10 to 20 carbon  
5 atoms. The fatty acids may be saturated or unsaturated and include,  
6 for example, lauric acid, myristic acid, stearic acid, oleic acid, linoleic acid and  
7 palmitic acid. Monoglyceride permeation enhancers include glycerol  
8 monooleate, glycerol monolaurate and glycerol monolinoleate, for example.  
9 In a presently preferred embodiment of this invention, the second permeation  
10 enhancer is a monoglyceride or a mixture of monoglycerides of unsaturated  
11 fatty acids, and more preferred is glycerol monolaurate (GML). As used  
12 herein and in the appended claims, the term "monoglyceride" refers to a  
13 monoglyceride or a mixture of monoglycerides of fatty acids.

14 It has been seen that glycerol monooleate having a total monoesters  
15 content of less than about 65% interacts adversely with known adhesive  
16 materials to such an extent that the adhesive cannot function to maintain a  
17 delivery device on the skin. Therefore, when an in-line adhesive is present as  
18 a part of the device of the invention so that a permeation enhancer must pass  
19 through the adhesive, and when glycerol monooleate is utilized as the second  
20 permeation enhancer, the glycerol monooleate must have a total monoesters  
21 content of at least 65%.

22 The permeation-enhancing mixture is dispersed throughout the matrix  
23 or carrier, preferably at a concentration sufficient to provide permeation-  
24 enhancing amounts of enhancer in the reservoir throughout the anticipated  
25 administration period. Where there is an additional, separate permeation  
26 enhancer matrix layer as well, as in FIG. 2, the permeation enhancer normally  
27 is present in the separate reservoir in excess of saturation.

28 One embodiment of a transdermal delivery device of the present  
29 invention is illustrated in FIG. 1. In FIG. 1, device 1 is comprised of a drug-  
30 and acyl lactylate-containing reservoir ("drug reservoir") 2 which is preferably

1 in the form of a matrix containing the drug and the enhancer dispersed  
2 therein. In those embodiments which include a mixture of permeation  
3 enhancers, drug reservoir 2 also includes these additional enhancers.  
4 A backing layer 3 is provided adjacent one surface of drug reservoir 2.  
5 Adhesive overlay 4 maintains the device 1 on the skin and may be fabricated  
6 together with, or provided separately from, the remaining elements of the  
7 device. With certain formulations, the adhesive overlay 4 may be preferable  
8 to an in-line contact adhesive, such as adhesive layer 28 as shown in FIG. 3.  
9 Backing layer 3 is preferably slightly larger than drug reservoir 2, and in this  
10 manner prevents the materials in drug reservoir 2 from adversely interacting  
11 with the adhesive in overlay 4. A strippable or removable liner 5 is also  
12 provided with device 1 and is removed just prior to application of device 1 to  
13 the skin.

14 Figure 2 illustrates another embodiment of the invention, device 10,  
15 shown in placement on the skin 17. In this embodiment, the transdermal  
16 delivery device 10 comprises a multi-laminate drug formulation/enhancer  
17 reservoir 11 having at least two zones 12 and 14. Zone 12 consists of a drug  
18 reservoir substantially as described with respect to FIG. 1. Zone 14  
19 comprises a permeation enhancer reservoir which is preferably made from  
20 substantially the same matrix as is used to form zone 12. Zone 14 comprises  
21 an acyl lactylate dispersed throughout, preferably in excess of saturation, and  
22 is substantially free of any undissolved drug. One or more additional  
23 permeation enhancers may optionally be included in zone 14 as well.  
24 A rate-controlling membrane 13 for controlling the release rate of the acyl  
25 lactylate and, optionally, any additional enhancers from zone 14 to zone 12 is  
26 placed between the two zones. A rate-controlling membrane (not shown) for  
27 controlling the release rate of the enhancer from zone 12 to the skin may also  
28 optionally be utilized and would be present between the skin 17 and zone 12.

1           The rate-controlling membrane may be fabricated from permeable,  
2 semipermeable or microporous materials which are known in the art to control  
3 the rate of agents into and out of delivery devices and having a permeability  
4 to the permeation enhancer lower than that of zone 12. Suitable materials  
5 include, but are not limited to, polyethylene, polyvinyl acetate and ethylene  
6 vinyl acetate copolymers.

7           An advantage of the device described in FIG. 2 is that the drug-loaded  
8 zone 12 is concentrated at the skin surface rather than throughout the entire  
9 mass of the reservoir 11. This functions to reduce the amount of drug in the  
10 device while maintaining an adequate supply of the permeation enhancer  
11 or mixture.

12           Superimposed over the drug formulation/enhancer -reservoir 11 of  
13 device 10 is a backing 15 and an adhesive overlay 16 as described above  
14 with respect to FIG. 1. In addition, a strippable liner (not shown) would  
15 preferably be provided on the device prior to use as described with respect to  
16 FIG. 1 and removed prior to application of the device 10 to the skin 17.

17           In the embodiments of FIGS. 1 and 2, the carrier or matrix material has  
18 sufficient viscosity to maintain its shape without oozing or flowing.  
19 If, however, the matrix or carrier is a low viscosity flowable material,  
20 the composition can be fully enclosed in a dense non-porous or microporous  
21 skin-contacting membrane, as known to the art from U.S. Pat. No. 4,379,454  
22 (noted above), for example.

23           An example of a presently preferred transdermal delivery device is  
24 illustrated in FIG. 3. In FIG. 3, transdermal delivery device 20 comprises a  
25 drug reservoir 22 containing together the drug and the acyl lactylate  
26 permeation enhancer. Optionally, one or more additional permeation  
27 enhancers may also be included in drug reservoir 22. Reservoir 22 is  
28 preferably in the form of a matrix containing the drug and the enhancer  
29 dispersed therein. Reservoir 22 is sandwiched between a backing layer 24,  
30 which is impermeable to both the drug and the acyl lactylate, and an in-line

1 contact adhesive layer 28. In FIG. 3, the drug reservoir 22 is formed of a  
2 material, such as a rubbery polymer, that is sufficiently viscous to maintain its  
3 shape. The device 20 adheres to the surface of the skin 17 by means of the  
4 contact adhesive layer 28. The adhesive for layer 28 should be chosen so  
5 that it is compatible and does not interact with any of the drug or, in particular,  
6 the acyl lactylate permeation enhancer. The adhesive layer 28 may optionally  
7 contain permeation enhancer and/or drug. A strippable liner (not shown) is  
8 normally provided along the exposed surface of adhesive layer 28 and is  
9 removed prior to application of device 20 to the skin 17. In an alternative  
10 embodiment, a rate-controlling membrane (not shown) is present and the  
11 drug reservoir 22 is sandwiched between backing layer 24 and the rate-  
12 controlling membrane, with adhesive layer 28 present on the skin-side of the  
13 rate-controlling membrane.

14 Various materials suited for the fabrication of the various layers of the  
15 transdermal devices of FIGS. 1, 2 or 3 are known in the art or are disclosed in  
16 the aforementioned transdermal device patents previously incorporated  
17 herein by reference.

18 The matrix making up the drug/acyl lactylate permeation enhancer  
19 reservoir can be a gel or a polymer. Suitable materials should be compatible  
20 with the drug and enhancer and any other components in the system.  
21 The matrix may be aqueous or non-aqueous based. Aqueous formulations  
22 typically comprise water or water/ethanol and about 1-5 wt% of a gelling  
23 agent, an example being a hydrophilic polymer such as hydroxyethylcellulose  
24 or hydroxypropylcellulose. Typical non-aqueous gels are comprised of  
25 silicone fluid or mineral oil. Mineral oil based gels also typically contain 1-2  
26 wt% of a gelling agent such as colloidal silicon dioxide. The suitability of a  
27 particular gel depends upon the compatibility of its constituents with the drug  
28 and the permeation-enhancing mixture in addition to any other components in  
29 the formulation.

1           When using a non-aqueous based formulation, the reservoir matrix is  
2 preferably composed of a hydrophobic polymer. Suitable polymeric matrices  
3 are well known in the transdermal drug delivery art, and examples are listed  
4 in the above-named patents previously incorporated herein by reference.  
5 A typical laminated system would consist essentially of a polymeric  
6 membrane and/or matrix such as ethylene vinyl acetate (EVA) copolymers,  
7 such as those described in US Patent No. 4,144,317, preferably having a vinyl  
8 acetate content in the range of from about 9% to about 60% and more  
9 preferably about 9% to 40% vinyl acetate. Polyisobutylene/oil polymers  
10 containing from 4-25% high molecular weight polyisobutylene and 20-81%  
11 low molecular weight polyisobutylene with the balance being an oil such as  
12 mineral oil or polyisobutylenes may also be used as the matrix material.

13           In addition to a drug and acyl lactylate, which are essential to the  
14 invention, the matrix may also contain stabilizers, dyes, pigments, inert fillers,  
15 tackifiers, excipients and other conventional components of transdermal  
16 delivery devices as are known in the art.

17           The amounts of the drug that are present in the therapeutic device,  
18 and that are required to achieve a therapeutic effect, depend on many  
19 factors, such as the minimum necessary dosage of the particular drug;  
20 the permeability of the matrix, of the adhesive layer and of the rate-controlling  
21 membrane, if present; and the period of time for which the device will be fixed  
22 to the skin. There is, in fact, no upper limit to the maximum amounts of drug  
23 present in the device. The minimum amount of each drug is determined by  
24 the requirement that sufficient quantities of drug must be present in the device  
25 to maintain the desired rate of release over the given period of application.

26           The drug is generally dispersed through the matrix at a concentration  
27 in excess of saturation, i.e. at unit activity. The amount of excess is  
28 determined by the intended useful life of the system. However, the drug may  
29 be present at initial levels below saturation without departing from this  
30 invention. Generally, the drug may be present at initially subsaturated levels

1 when: 1) the skin flux of the drug is sufficiently low such that the reservoir  
2 drug depletion is slow and small; 2) non-constant delivery of the drug is  
3 desired or acceptable; and/or 3) saturation of the reservoir is achieved in use  
4 due to migration of water into the reservoir from the skin, where water is  
5 abundantly available.

6 The acyl lactylate permeation enhancer is dispersed throughout the  
7 matrix, preferably at a concentration sufficient to provide permeation--  
8 enhancing concentrations of enhancer in the reservoir throughout the  
9 anticipated administration period.

10 In certain embodiments of the invention, one or more additional  
11 permeation enhancers, such as a monoglyceride or mixture of  
12 monoglycerides of fatty acids including glycerol monolaurate (GML) and  
13 glycerol monooleate (GMO), lauramide diethanolamine (LDEA), esters of fatty  
14 acids having from about 10 to 20 carbon atoms, and/or a lower C<sub>1-4</sub> alcohol  
15 such as ethanol or isopropanol, may also be dispersed throughout the matrix,  
16 preferably at a concentration sufficient to provide permeation-enhancing  
17 concentrations of enhancer in the drug reservoir throughout the anticipated  
18 administration period.

19 In the present invention, the drug is delivered through the skin or other  
20 body surface at a therapeutically effective rate (that is, a rate that provides an  
21 effective therapeutic result) and the acyl lactylate permeation enhancer is  
22 delivered at a permeation-enhancing rate (that is, a rate that provides  
23 increased permeability of the application site to the drug) for a predetermined  
24 time period.

25 A preferred embodiment of the present invention is a monolith such as  
26 that illustrated in FIG. 3 (either with or without a rate-controlling membrane)  
27 wherein reservoir 22 comprises, by weight, 30- 90% polymer (preferably EVA  
28 with a vinyl acetate content of 40%), 0.01-40% drug, and 1-70% of an acyl  
29 lactylate. The in-line adhesive layer 28 contains an adhesive which is  
30 compatible with the permeation enhancer. In another preferred embodiment

1 of the invention, a monolith such as that in FIG. 3 includes reservoir 22  
2 comprising, by weight, 30-90% polymer (preferably EVA with a vinyl acetate  
3 content of 40%), 0.01-40% drug, 1-70% acyl lactylate, and 1-45% of a second  
4 permeation enhancer, preferably GML.

5 The devices of this invention can be designed to effectively deliver a  
6 drug for an extended time period of up to 7 days or longer. Seven days is  
7 generally the maximum time limit for application of a single device because  
8 the skin site may be affected by a period of occlusion greater than 7 days,  
9 or other problems such as the system or edges of the system lifting off of the  
10 skin may be encountered over such long periods of application. Where it is  
11 desired to have drug delivery for greater than 7 days (such as, for example,  
12 when a hormone is being applied for a contraceptive effect), when one device  
13 has been in place on the skin for its effective time period, it is replaced with a  
14 fresh device, preferably on a different skin site.

15 The transdermal therapeutic devices of the present invention are  
16 prepared in a manner known in the art, such as by those procedures,  
17 for example, described in the transdermal device patents listed previously  
18 herein. The following examples are offered to illustrate the practice of the  
19 present invention and are not intended to limit the invention in any manner.

#### 20 21 EXAMPLE 1

22  
23 Several test samples were made to measure the progesterone flux  
24 through human cadaver skin from donor vehicles containing progesterone at  
25 saturation in mineral oil. The progesterone vehicle was also mixed with 20%  
26 by weight caproyl lactic acid or lauroyl lactic acid, or dispersed with 1.1%  
27 by weight glycerol monolaurate. Transdermal fluxes were obtained using  
28 human epidermis at 35° C in standard diffusion cells. As seen in Figure 4,  
29 the progesterone mixtures with caproyl lactic acid and lauroyl lactic acid  
30 demonstrated superior flux of progesterone through skin.

## EXAMPLE 2

Several test samples were made to measure the testosterone flux through human cadaver skin. Testosterone saturated in mineral oil was used as a control, and was compared with mixtures including 12% GML / 7% caproyl lactic acid, or 12% GML / 7% lauroyl lactic acid. Transdermal fluxes were obtained using human epidermis at 35° C in standard diffusion cells. As demonstrated in Figure 5, the solutions containing the GML/acyl lactate mixtures resulted in a testosterone flux through skin at least double that of the control.

## EXAMPLE 3

The drug/permeation enhancer reservoirs were prepared by mixing ethylene vinyl acetate having a vinyl acetate content of 40 percent ("EVA 40", USI Chemicals, Illinois) in an internal mixer (Brabender type mixer) until the EVA 40 pellets fused. Testosterone, lauroyl lactic acid (LLA), caproyl lactic acid (CLA), GML, lactic acid (LA), M-DEA, L-DEA, or lauryl lactate (LL) were then added as shown in Table 1. The mixture was blended, cooled and calendered to a 5 mil thick film.

The film was then laminated to a Medpar® (3M) backing on one side and an acrylate contact adhesive (3M) on the opposite side. The laminate was then cut into 1.98 cm<sup>2</sup> circles using a steel punch.

Circular pieces of human epidermis were mounted on horizontal permeation cells with the stratum corneum facing the donor compartment of the cell. The release liner of the laminate was removed and the systems were centered over the stratum corneum side of the epidermis. The cells were then masked. A known volume of receptor solution (20 ml) was

equilibrated at 35 °C and placed in the receptor compartment. Air bubbles were removed from the receptor compartment, the cell was capped and placed in a water bath shaker at 35 °C.

Table 1

Drug/ Permeation Enhancer Reservoir (wt %)

RESERVOIR FORMULATION	WEIGHT PERCENT
Testosterone/LLA/GML/EVA 40	10/20/20/50
Testosterone/CLA/ L-DEA/EVA 40	10/20/20/50
Testosterone/LL/LA/GML/EVA 40	10/12/3/20/55
Testosterone/LL/LA/L-DEA/EVA 40	10/12/3/20/55
Testosterone/LLA/M-DEA/EVA 40	10/20/20/50
Testosterone/LL/LA/M-DEA/EVA 40	10/12/3/20/55
Testosterone/EVA 40	2/98

At given time intervals, the entire receptor solution was removed from the cells and replaced with an equal volume of fresh receptor solutions previously equilibrated at 35 °C. The receptor solutions are stored in capped vials at room temperature until assayed for testosterone content by HPLC. From the drug concentration and the volume of the receptor solutions, the area of permeation and the time interval, the flux of the drug through the epidermis was calculated as follows: (drug concentration X volume of receptor)/( area X time) = flux ( $\mu\text{g}/\text{cm}^2 \cdot \text{hr}$ ). The flux of the testosterone achieved from the various systems is shown in Figure 6. As demonstrated in Figure 6, formulations comprising mixtures including lauroyl lactic acid or caproyl lactic acid resulted in superior flux of testosterone through the skin.

- 1 This invention has been described in detail with particular reference to certain
- 2 preferred embodiments thereof, but it will be understood that variations and
- 3 modifications can be effected within the spirit and scope of
- 4 the invention.

1   What is claimed is:

2

3           1.     A composition of matter for transdermally delivering a drug to  
4   the systemic circulation of a patient by permeation through a body surface or  
5   membrane comprising, in combination:

6           (a) a drug; and

7           (b) a permeation-enhancing amount of an acyl lactylate permeation  
8   enhancer sufficient to substantially increase the permeability of the body  
9   surface or membrane to the drug in order to systemically deliver said drug to  
10   a patient at a therapeutically effective rate, wherein the drug and acyl lactylate  
11   permeation enhancer are dispersed within a pharmaceutically acceptable  
12   carrier.

13          2.     A composition according to claim 1 wherein the acyl lactylate is  
14   caproyl lactic acid.

15          3.     A composition according to claim 1 wherein the acyl lactylate is  
16   lauroyl lactic acid.

17          4.     A composition according to claim 1 wherein the acyl lactylate  
18   permeation enhancer is combined with a permeation-enhancing amount of  
19   one or more of the permeation enhancers selected from the group consisting  
20   of monoglycerides or mixtures of monoglycerides of fatty acids, lauramide  
21   diethanolamine, esters of fatty acids having from about 10 to 20 carbon  
22   atoms, and lower C<sub>1-4</sub> alcohols.

23          5.     A composition according to claim 1 wherein the acyl lactylate  
24   permeation enhancer is combined with a permeation-enhancing amount of  
25   glycerol monolaurate.

1           6.     A device (1,20) for the transdermal administration of a drug to  
2     the systemic circulation of a patient at a therapeutically effective rate, by  
3     permeation through a body surface or membrane, comprising:

4           (a) a reservoir (2,22) comprising a drug and a permeation enhancing-  
5     amount of an acyl lactylate permeation enhancer ; and

6           (b) means for maintaining (4,28) said reservoir in drug- and  
7     permeation enhancer- transmitting relation with the body surface or  
8     membrane, wherein said drug is delivered to the systemic circulation of a  
9     patient at a therapeutically effective rate.

10          7.     A device according to claim 6 wherein the acyl lactylate is  
11     caproyl lactic acid.

12          8.     A device according to claim 6 wherein the acyl lactylate is  
13     lauroyl lactic acid.

14          9.     A device according to claim 6 wherein the drug is testosterone.

15          10.    A device according to claim 6 wherein the drug is progesterone.

16          11.    A device according to claim 6 wherein the acyl lactylate  
17     permeation enhancer is combined with a permeation-enhancing amount of  
18     one or more of the permeation enhancers selected from the group consisting  
19     of monoglycerides or mixtures of monoglycerides of fatty acids, lauramide  
20     diethanolamine, esters of fatty acids having from about 10 to 20 carbon  
21     atoms, and lower C<sub>1-4</sub> alcohols.

22          12.    A device (10) for the transdermal administration of a drug to the  
23     systemic circulation of a patient at a therapeutically effective rate,  
24     by permeation through a body surface or membrane, comprising:

25           (a) a first reservoir (12) comprising a drug and a permeation-  
26     enhancing amount of an acyl lactylate permeation enhancer;

27           (b) a second reservoir (14) comprising an additional amount of said  
28     permeation enhancer and substantially free of said drug;

29           (c) a rate controlling membrane (13) between the first reservoir and  
30     the second reservoir;

1           (d) means for maintaining (16) said first and second reservoirs in drug-  
2 and permeation enhancer- transmitting relation with the body surface or  
3 membrane, wherein the drug is delivered to the systemic circulation of a  
4 patient at a therapeutically effective rate.

5           13. A device according to claim 10 wherein the acyl lactylate is  
6 caproyl lactic acid.

7           14. A device according to claim 10 wherein the acyl lactylate is  
8 lauroyl lactic acid.

9           15. A device according to claim 10 wherein the drug is testosterone.

10          16. A device according to claim 10 wherein the drug is  
11 progesterone.

12          17. A device according to claim 10 wherein the acyl lactylate  
13 permeation enhancer is combined with a permeation-enhancing amount of  
14 one or more of the permeation enhancers selected from the group consisting  
15 of monoglycerides or mixtures of monoglycerides of fatty acids, lauramide  
16 diethanolamine, esters of fatty acids having from about 10 to 20 carbon  
17 atoms, and lower C<sub>1-4</sub> alcohols.

18          18. A method for the transdermal administration of a drug to the  
19 systemic circulation of a patient, at a therapeutically effective rate, by  
20 permeation through a body surface or membrane, comprising:

21           (a) simultaneously administering, to the body surface or membrane,  
22 a drug; and

23           (b) a permeation-enhancing amount of an acyl lactylate permeation  
24 enhancer sufficient to substantially increase the permeability of the body  
25 surface or membrane to the drug in order to systemically deliver said drug to  
26 a patient at a therapeutically effective rate.

1           19.    A method according to claim 18 further comprising  
2 simultaneously coadministering a permeation-enhancing amount of one or  
3 more of the permeation enhancers selected from the group consisting of  
4 monoglycerides or mixtures of monoglycerides of fatty acids, lauramide  
5 diethanolamine, esters of fatty acids having from about 10 to 20 carbon  
6 atoms, and lower C<sub>1-4</sub> alcohols.

7           20.    A method according to claim 18 wherein the acyl lactylate is  
8 lauroyl lactic acid.

9           21.    A method according to claim 18 wherein the acyl lactylate is  
10 caproyl lactic acid.

11          22.    A method according to claim 18 wherein the drug is  
12 testosterone.

13          23.    A method according to claim 18 wherein the drug is  
14 progesterone.

15          24.    A method according to claim 18 wherein the drug is  
16 administered to the systemic circulation of said patient at a therapeutically  
17 effective rate throughout a substantial portion of an administration period of at  
18 least about 10 hours.

19          25.    A method for the transdermal administration of a drug, the  
20 method comprising the step of placing a transdermal delivery device (1,20)  
21 onto a body surface or membrane of a person, the transdermal delivery  
22 device comprising:

23           (a) a reservoir (2,22) comprising a drug and a permeation-enhancing  
24 amount of an acyl lactylate permeation enhancer; and

25           (b) means for maintaining (4,28) said reservoir in drug- and  
26 permeation enhancer- transmitting relation with the body surface or  
27 membrane, wherein the drug is delivered to the systemic circulation of a  
28 patient at a therapeutically effective rate.

- 1           26.    A method for the transdermal administration of a drug, the  
2 method comprising the step of placing a transdermal delivery device (10)  
3 onto a body surface or membrane of a person, the transdermal delivery  
4 device comprising:
- 5           (a) a first reservoir (10) comprising a drug and a permeation-  
6 enhancing amount of an acyl lactylate permeation enhancer; and
- 7           (b) a second reservoir (14) comprising an additional amount of said  
8 permeation enhancer and substantially free of said drug;
- 9           (c) a rate controlling membrane (13) between the first reservoir and  
10 the second reservoir;
- 11          (d) means for maintaining (16) said first and second reservoirs in drug-  
12 and permeation enhancer- transmitting relation with the body surface or  
13 membrane, wherein the drug is delivered to the systemic circulation of a  
14 patient at a therapeutically effective rate.




FIG. 1




FIG. 2






FIG. 3




- CONTROL: PROGESTERONE SATURATED IN M.O.  
● CAPROYL LACTYLIC ACID 20% IN M.O.  
▽ LAUROYL LACTYLIC ACID 20% IN M.O.  
▼ GML 1.1%(DISPERSED) IN M.O.

FIG. 4



○ CONTROL: TESTOSTERONE SATURATED IN M.O.  
● GML 12%/CAPROYL LACTYLIC ACID 7% IN M.O.  
▽ GML 12%/LAUROYL LACTYLIC ACID 7% IN M.O.

FIG. 5



- FORMULATION 1
- CONTROL WITHOUT ADHESIVE
- ▽ FORMULATION 4
- ▼ FORMULATION 6
- FORMULATION 8
- FORMULATION 10
- △ FORMULATION 12

N=7 (4 SKIN DONORS)

FIG. 6a

## FORMULATION 1

LAUROYL LACTIC ACID	20%
GML	20%
EVA40%va	50%
TESTOSTERONE	10%

## FORMULATION 4

CAPROYL LACTIC ACID	20%
L-DEA	20%
EVA40%va	50%
TESTOSTERONE	10%

## FORMULATION 6

LAURYL LACTATE	12%
LACTIC ACID	3%
GML	20%
EVA40%va	55%
TESTOSTERONE	10%

## FORMULATION 8

LAURYL LACTATE	12%
LACTIC ACID	3%
L-DEA	20%
EVA40%va	55%
TESTOSTERONE	10%

## FORMULATION 10

LAUROYL LACTIC ACID	20%
M-DEA	20%
EVA40%va	50%
TESTOSTERONE	10%

## FORMULATION 12

LAUROYL LACTIC	12%
LACTIC ACID	3%
M-DEA	20%
EVA40%va	55%
TESTOSTERONE	10%

## CONTROL

EVA40%va	98%
TESTOSTERONE	2%

FIG. 6b

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K47/14 A61L15/44

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,5 344 651 (SCHWEN R. J. ET AL) 6 September 1994 see column 5, line 18 - line 22 see column 12; examples 4,6 ---	1
X	WO,A,88 06880 (R.I.T.A. CORPORATION) 22 September 1988 see page 6, line 1 - line 9 see page 7, line 1 - line 11 see page 5, line 1 - line 11 ---	1
X	US,A,5 124 354 (GREEN M. R.) 23 June 1992 cited in the application see column 16, line 5 - line 7 see column 16, line 23 --- -/-	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

2 September 1996

Date of mailing of the international search report

13. 09. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Boulois, D

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,95 05153 (UNILEVER PLC) 23 February 1995 see page 5, line 23 - line 25 ---	1
A	WO,A,95 09006 (ALZA CORPORATION) 6 April 1995 see claims 1,12,14,16,17 -----	1

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 06868

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 18-26 are directed to a method of treatment of the human body, the search has been carried out and based on the alleged effects of the composition.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5344651	06-09-94	WO-A- 9503319	02-02-95
WO-A-8806880	22-09-88	CA-A- 1329364	10-05-94
		US-A- 4822601	18-04-89
		WO-A- 9101713	21-02-91
		CA-A- 1316829	27-04-93
		DE-A- 3882848	09-09-93
		DE-T- 3882848	13-01-94
		EP-A- 0305493	08-03-89
		JP-A- 7173045	11-07-95
		JP-B- 7068148	26-07-95
		JP-T- 2500358	08-02-90
		US-A- 4946832	07-08-90
US-A-5124354	23-06-92	CA-A- 2018737	14-12-90
		EP-A- 0403238	19-12-90
		JP-A- 3063213	19-03-91
WO-A-9505153	23-02-95	AU-B- 7499294	14-03-95
		AU-B- 7612194	14-03-95
		CA-A- 2166468	23-02-95
		CA-A- 2166469	23-02-95
		WO-A- 9505160	23-02-95
		EP-A- 0713385	29-05-96
		ZA-A- 9406128	17-07-95
WO-A-9509006	06-04-95	AU-B- 7924994	18-04-95
		AU-B- 7964794	18-04-95
		CA-A- 2165802	06-04-95
		CA-A- 2167526	06-04-95
		EP-A- 0721348	17-07-96
		EP-A- 0721349	17-07-96
		WO-A- 9509007	06-04-95